

Course References

Lloyd-Williams M, Shiels C, Taylor F, Dennis M. Depression - An independent predictor of early death in patients with advanced Cancer. *Journal of affective disorders*. 2008;113:127-132. doi:[10.1016/j.jad.2008.04.002](https://doi.org/10.1016/j.jad.2008.04.002)

Voogd JN de, Wempe JB, Koëter GH, et al. Depressive Symptoms as predictors of mortality in patients with COPD. *CHEST*. 2009;135(3):619-625. doi:[10.1378/chest.08-0078](https://doi.org/10.1378/chest.08-0078)

Katon WJ. Epidemiology and treatment of depression in patients with chronic medical illness. *Dialogues Clin Neurosci*. 2011;13(1):7-23. doi:[10.31887/DCNS.2011.13.1/wkaton](https://doi.org/10.31887/DCNS.2011.13.1/wkaton)

Tara Smedley, PharmD, BCPS, APh. How Long does it take for antidepressants to start working? GoodRx. Published August 10, 2022. Accessed February 20, 2023. <https://www.goodrx.com/healthcare-access/medication-education/how-long-does-it-take-for-antidepressants-to-work>

Mental Health Disparities: Diverse Populations. Accessed February 17, 2023. <https://www.psychiatry.org/443/psychiatrists/diversity/education/mental-health-facts>

Watanabe N, Omori IM, Nakagawa A, et al. Mirtazapine versus other antidepressive agents for depression. *Cochrane Database of Systematic Reviews*. 2011;(12). doi:[10.1002/14651858.CD006528.pub2](https://doi.org/10.1002/14651858.CD006528.pub2)

Patient Race/Ethnicity and Quality of Patient–Physician Communication During Medical Visits. *AJPH*. Vol. 94 Issue 12. Accessed February 17, 2023. <https://ajph.aphapublications.org/doi/full/10.2105/AJPH.94.12.2084>

Relationship between depression and exacerbations in COPD. European Respiratory Society. Accessed February 17, 2023. <https://erj.ersjournals.com/content/32/1/53>

Clarke TC, Barnes PM, Black LI, Stussman BJ, Nahin RL. Use of yoga, meditation, and chiropractors among U.S. adults aged 18 and over. *NCHS Data Brief*. 2018;(325):1-8. Accessed February 16, 2023. <https://www.cdc.gov/nchs/data/databriefs/db325-h.pdf>.

Clarke TC, Black LI, Stussman BJ, Barnes PM, Nahin RL. Trends in the use of complementary health approaches among adults: United States, 2002-2012. *Natl Health Stat Report*. 2015;(79):1-16. Accessed February 15, 2023. <https://pubmed.ncbi.nlm.nih.gov/25671660/>.

Additional Resources

American Psychiatric Association. Mental Health Disparities: Diverse Populations. Accessed March 7, 2023. <https://www.psychiatry.org/psychiatrists/cultural-competency/education/mental-health-facts>

Blazer DG, Kessler RC, McGonagle KA, Swartz MS. The prevalence and distribution of major depression in a national community sample: the National Comorbidity Survey. *Am J Psychiatry*. 1994;151(7):979-986. doi:[10.1176/ajp.151.7.979](https://doi.org/10.1176/ajp.151.7.979)

Block SD. Assessing and managing depression in the terminally ill patient. ACP-ASIM End-of-Life Care Consensus Panel. American College of Physicians - American Society of Internal Medicine. *Ann Intern Med*. 2000;132(3):209-218. doi:[10.7326/0003-4819-132-3-200002010-00007](https://doi.org/10.7326/0003-4819-132-3-200002010-00007)

Block SD. Psychological issues in end-of-life care. *J Palliat Med*. 2006;9(3):751-772. doi:[10.1089/jpm.2006.9.751](https://doi.org/10.1089/jpm.2006.9.751)

Breitbart W, Bruera E, Chochinov H, Lynch M. Neuropsychiatric syndromes and psychological symptoms in patients with advanced cancer. *J Pain Symptom Manage*. 1995;10(2):131-141. doi:[10.1016/0885-3924\(94\)00075-v](https://doi.org/10.1016/0885-3924(94)00075-v)

Chochinov HM, Hack T, Hassard T, Kristjanson LJ, McClement S, Harlos M. Dignity therapy: a novel psychotherapeutic intervention for patients near the end of life. *J Clin Oncol*. 2005;23(24):5520-5525. doi:[10.1200/JCO.2005.08.391](https://doi.org/10.1200/JCO.2005.08.391)

Chochinov HM, Kristjanson LJ, Breitbart W, et al. Effect of dignity therapy on distress and end-of-life experience in terminally ill patients: a randomised controlled trial. *Lancet Oncol*. 2011;12(8):753-762. doi:[10.1016/S1470-2045\(11\)70153-X](https://doi.org/10.1016/S1470-2045(11)70153-X)

Chochinov HM, Tatarzyn D, Clinch JJ, Dudgeon D. Will to live in the terminally ill. *Lancet*. 1999;354(9181):816-819. doi:[10.1016/S0140-6736\(99\)80011-7](https://doi.org/10.1016/S0140-6736(99)80011-7)

Christakis NA. Timing of referral of terminally ill patients to an outpatient hospice. *J Gen Intern Med*. 1994;9(6):314-320. doi:[10.1007/BF02599178](https://doi.org/10.1007/BF02599178)

Consumer Reports Best Buy Drugs. Using Anti-Depressants to Treat Depression: Comparing Effectiveness, Safety and Price. Consumer Reports Website. Updated 2013. Accessed March 7, 2023. <https://consumerhealthchoices.org/wp-content/uploads/2012/08/BBD-Antidepressant-Full.pdf>

Dworkin RH, O'Connor AB, Audette J, et al. Recommendations for the pharmacological management of neuropathic pain: an overview and literature update. *Mayo Clin Proc*. 2010;85(3 Suppl):S3-S14. doi:[10.4065/mcp.2009.0649](https://doi.org/10.4065/mcp.2009.0649)

Geschwind N, Peeters F, Huibers M, van Os J, Wichers M. Efficacy of mindfulness-based cognitive therapy in relation to prior history of depression: randomised controlled trial. *Br J Psychiatry*. 2012;201(4):320-325. doi:[10.1192/bjp.bp.111.104851](https://doi.org/10.1192/bjp.bp.111.104851)

Goodwin RD, Marusic A, Hoven CW. Suicide attempts in the United States: the role of physical illness. *Soc Sci Med*. 2003;56(8):1783-1788. doi:[10.1016/s0277-9536\(02\)00174-0](https://doi.org/10.1016/s0277-9536(02)00174-0)

Hinton J. Which patients with terminal cancer are admitted from home care?. *Palliat Med*. 1994;8(3):197-210. doi:[10.1177/026921639400800303](https://doi.org/10.1177/026921639400800303)

Jacobs S. Pathologic Grief: Maladaptation to Loss. Washington, DC: American Psychiatric Press, Inc., 1993.

Kessler RC, Borges G, Walters EE. Prevalence of and risk factors for lifetime suicide attempts in the National Comorbidity Survey. *Arch Gen Psychiatry*. 1999;56(7):617-626. doi:[10.1001/archpsyc.56.7.617](https://doi.org/10.1001/archpsyc.56.7.617)

Kroenke K, Spitzer RL, Williams JB, Löwe B. An ultra-brief screening scale for anxiety and depression: the PHQ-4. *Psychosomatics*. 2009;50(6):613-621. doi:[10.1176/appi.psy.50.6.613](https://doi.org/10.1176/appi.psy.50.6.613)

Lee YC, Chen PP. A review of SSRIs and SNRIs in neuropathic pain. *Expert Opin Pharmacother*. 2010;11(17):2813-2825. doi:[10.1517/14656566.2010.507192](https://doi.org/10.1517/14656566.2010.507192)

Lloyd-Williams M, Friedman T, Rudd N. A survey of antidepressant prescribing in the terminally ill. *Palliat Med*. 1999;13(3):243-248. doi:[10.1191/026921699676753309](https://doi.org/10.1191/026921699676753309)

Malhi GS, Byrow Y, Bassett D, et al. Stimulants for depression: On the up and up?. *Aust N Z J Psychiatry*. 2016;50(3):203-207. doi:[10.1177/0004867416634208](https://doi.org/10.1177/0004867416634208)

Marks DM, Shah MJ, Patkar AA, Masand PS, Park GY, Pae CU. Serotonin-norepinephrine reuptake inhibitors for pain control: premise and promise. *Curr Neuropharmacol*. 2009;7(4):331-336. doi:[10.2174/157015909790031201](https://doi.org/10.2174/157015909790031201)

Meyer HA, Sinnott C, Seed PT. Depressive symptoms in advanced cancer. Part 2. Depression over time; the role of the palliative care professional. *Palliat Med*. 2003;17(7):604-607. doi:[10.1191/0269216303pm813oa](https://doi.org/10.1191/0269216303pm813oa)

Montross L, Winters KD, Irwin SA. Dignity therapy implementation in a community-based hospice setting. *J Palliat Med*. 2011;14(6):729-734. doi:[10.1089/jpm.2010.0449](https://doi.org/10.1089/jpm.2010.0449)

National Center for Complementary and Integrative Health, National Institutes of Health: Complementary, Alternative, or Integrative Health: What's In a Name? Updated April 2021. Accessed March 7, 2023. <https://www.nccih.nih.gov/health/complementary-alternative-or-integrative-health-whats-in-a-name>

National Center for Complementary and Integrative Health, National Institutes of Health: Cancer: In Depth. Updated October 2021. Accessed March 7, 2023. <https://www.nccih.nih.gov/health/cancer-and-complementary-health-approaches-what-you-need-to-know>

O'Mahony S, Goulet J, Kornblith A, et al. Desire for hastened death, cancer pain and depression: report of a longitudinal observational study. *J Pain Symptom Manage*. 2005;29(5):446-457. doi:[10.1016/j.jpainsymman.2004.08.010](https://doi.org/10.1016/j.jpainsymman.2004.08.010)

Passik SD, Dugan W, McDonald MV, Rosenfeld B, Theobald DE, Edgerton S. Oncologists' recognition of depression in their patients with cancer. *J Clin Oncol*. 1998;16(4):1594-1600. doi:[10.1200/JCO.1998.16.4.1594](https://doi.org/10.1200/JCO.1998.16.4.1594)

Pelletier G, Verhoef MJ, Khatri N, Hagen N. Quality of life in brain tumor patients: the relative contributions of depression, fatigue, emotional distress, and existential issues. *J Neurooncol*. 2002;57(1):41-49. doi:[10.1023/a:1015728825642](https://doi.org/10.1023/a:1015728825642)

Portenoy RK, Thaler HT, Kornblith AB, et al. Symptom prevalence, characteristics and distress in a cancer population. *Qual Life Res*. 1994;3(3):183-189. doi:[10.1007/BF00435383](https://doi.org/10.1007/BF00435383)

Purgato M, Papola D, Gastaldon C, et al. Paroxetine versus other anti-depressive agents for depression. *Cochrane Database Syst Rev*. 2014;(4):CD006531. Published 2014 Apr 3. doi:[10.1002/14651858.CD006531.pub2](https://doi.org/10.1002/14651858.CD006531.pub2)

Regier DA, Boyd JH, Burke JD Jr, et al. One-month prevalence of mental disorders in the United States. Based on five Epidemiologic Catchment Area sites. *Arch Gen Psychiatry*. 1988;45(11):977-986. doi:[10.1001/archpsyc.1988.01800350011002](https://doi.org/10.1001/archpsyc.1988.01800350011002)

Riordan P, Briscoe J, Kamal AH, Jones CA, Webb JA. Top Ten Tips Palliative Care Clinicians Should Know About Mental Health and Serious Illness. *J Palliat Med*. 2018;21(8):1171-1176. doi:[10.1089/jpm.2018.0207](https://doi.org/10.1089/jpm.2018.0207)

Saarto T, Wiffen PJ. Antidepressants for neuropathic pain. *Cochrane Database Syst Rev*. 2007;(4):CD005454. Published 2007 Oct 17. doi:[10.1002/14651858.CD005454.pub2](https://doi.org/10.1002/14651858.CD005454.pub2)

Seow H, Barbera L, Sutradhar R, et al. Trajectory of performance status and symptom scores for patients with cancer during the last six months of life. *J Clin Oncol*. 2011;29(9):1151-1158. doi:[10.1200/JCO.2010.30.7173](https://doi.org/10.1200/JCO.2010.30.7173)

Sharpe M, Strong V, Allen K, et al. Major depression in outpatients attending a regional cancer centre: screening and unmet treatment needs. *Br J Cancer*. 2004;90(2):314-320. doi:[10.1038/sj.bjc.6601578](https://doi.org/10.1038/sj.bjc.6601578)

Sohail Z, Bailey RK, Richie WD. Misconceptions of depression in african americans. *Front Psychiatry*. 2014;5:65. Published 2014 Jun 20. doi:[10.3389/fpsy.2014.00065](https://doi.org/10.3389/fpsy.2014.00065)

Wilson KG, Chochinov HM, McPherson CJ, et al. Suffering with advanced cancer. *J Clin Oncol*. 2007;25(13):1691-1697. doi:[10.1200/JCO.2006.08.6801](https://doi.org/10.1200/JCO.2006.08.6801)

Wilson KG, Chochinov HM, Skirko MG, et al. Depression and anxiety disorders in palliative cancer care. *J Pain Symptom Manage*. 2007;33(2):118-129. doi:[10.1016/j.jpainsymman.2006.07.016](https://doi.org/10.1016/j.jpainsymman.2006.07.016)

Updated: 2/20/2023